

HIV NURSING 2019



21-22 September 2019 • Rome, Italy

HOSTED BY:



IN PARTNERSHIP WITH:



‘Zero, No Risk, Can’t pass it On’

U=U: A Multidisciplinary Approach

Shaun Watson, Clinical Nurse Specialist (HIV Community)



HIV NURSING CONFERENCE | 21-22 September 2019 • Rome, Italy 

Declarations

I have undertaken paid work and received scholarships from

ViiV, Gilead, MSD & Pfizer.

Why are we discussing U=U?

- **Not everyone is onboard as messages differ from 'zero risk' 'virtually no risk' 'negligible' this can be confusing.**
- "I have a patient who..."
- **What about viral load is it >200 >50 >20?**
- "I just don't understand the science...."

The Evidence

Swiss Statement	Expert opinion and evidence review of >25 smaller studies looking at impact of ART on risk factors for HIV transmission.	Concluded that transmission would not occur undetectable with viral load.	2008	Vernazza P et al. [7]
HPTN 052	1763 serodifferent heterosexual couples randomised to immediate or deferred ART. Although condom use was high the impact of ART was highly significant.	All infections occurred in people with detectable viral load: n=17 in the deferred ART group and one early infection in the ART group before VL was undetectable. Follow-up reported out to four years.	2011	Cohen M et al. [8, 9]

“an HIV positive person on effective HIV treatment (ART) cannot transmit HIV through sexual contact”

- on ART and adherent
- undetectable VL
- no STIs
- risk <1 in 100,000 (<0.001%)

The Evidence

PARTNER	Prospective observational European study in ~900 serodifferent couples who were not using condoms.	Final results reported zero transmissions after more than 58,000 times couples had sex without condoms when viral load was undetectable <200 copies/mL.	2014 (interim). 2016 (final)	Rodgers A et al. [10, 11]
Opposites Attract	Prospective observational study in 358 serodifferent gay male couples in Australia, Thailand and Brazil.	Zero transmissions when viral load was undetectable <200 copies/mL.	2017	Grulich A et al. [12]
PARTNER2	Extension of PARTNER study to collect additional follow-up in gay male couples.	Study is fully recruited and still ongoing (2014–2017).	Expected 2018.	[13]

PARTNER 2

- **900 gay couples only (March 2014 to May 2018)**
- **Aim - to provide a similar level of confidence for gay men to PARTNER**
- **The PARTNER study recruited HIV couples (one partner positive, one negative) at 75 clinical sites in 14 European countries. They tested the HIV-negative partners every six to 12 months for HIV, and tested viral load in the HIV-positive partners. Both partners also completed behavioural surveys. In cases of HIV infection in the negative partners, their HIV was genetically analysed to see if it came from their regular partner.**
- **The study found no transmissions between gay couples where the HIV-positive partner had a viral load under 200 copies/ml – even though there were nearly 77,000 acts of condomless sex between them.**

Risk of HIV transmission through condomless sex in serodifferent gay couples with the HIV-positive partner taking suppressive antiretroviral therapy (PARTNER): final results of a multicentre, prospective, observational study. Rodger, Alison JColl, Pep et al (2019) The Lancet , Volume 0 , Issue 0

Partner 2 at AIDS2018

Q: What would you say to HIV information providers who are withholding this [U=U] information from people with HIV?

A: "It is very, very clear that the risk is zero. If you are on suppressive ART you are sexually non-infectious. *THE TIME FOR EXCUSES IS OVER.*"

**Dr. Alison Rodger, lead author of PARTNER2 at the International AIDS Conference
- AIDS 2018 [presentation](#) (July, 2018)**

This means...

In absence of evidence, having an opinion against U=U is either out-of-date or just prejudice - Simon Collins (i-Base)

- If you are HIV positive and undetectable you cannot pass the virus on to sexual partners.**
- The challenge since 2008 is whether HIV transmission can occur – so far this not been proven.**

There is no evidence, anywhere, that anyone has transmitted HIV when they have an undetectable viral load


It's a fact
U=U

Undetectable equals untransmittable



The British HIV Association is proud to support the #UequalsU consensus statement of the Prevention Access Campaign


The Campaign



U=U
#LIETEProject

UNDETECTABLE = UNTRANSMITTABLE

PRIDE for YOUTH
LIVING WITH HIV?
Taking HIV medication daily suppresses the virus and means you can't pass it to others.
Need help getting effective treatment for HIV? Pride for Youth can help! Call us today at (516) 679-9000 to get connected.
Visit us at www.pridefor youth.org.uk



nam aidsmap
HIV & AIDS - sharing knowledge, changing lives

"The scientific evidence is clear. Someone whose HIV is undetectable does not pose an infection risk to their sexual partners."

For information on HIV you can rely on: www.aidsmap.com #UequalsU

U=U Undetectable Equals Untransmittable




I CAN'T PASS ON HIV.
People on effective HIV treatment cannot pass on the virus.

Get the facts
StartsWithMe.org.uk

IT STARTS WITH ME
TERRENCE HIGGINS TRUST

Produced by Terrence Higgins Trust for HIV PREVENTION ENGLAND.
Terrence Higgins Trust is a registered charity in England and Wales (reg. no. 205527) and in Scotland (SC031996). Company reg. no. 1728492. Photography by IFCOM ENGLAND.



YOU are the solution

Taking HIV medication daily suppresses the virus and means you can't pass it to others.

#UequalsU
DCTakesonHIV.com



CAN'T PASS IT ON

People on effective HIV treatment **CANNOT** pass on the virus

TERRENCE HIGGINS TRUST

FACT

That's Amazing But....

- **This is not a campaign to promote unprotected sex.**
- **We still need studies on breastfeeding and blood transmission.**
- **There are people living with HIV who despite everything cannot get to undetectable.**
- **Globally there are people living with HIV who cannot access ART.**
- **There are people living with HIV who decide not to take treatment (intentional non-adherence)**

What Can We Do?

- It's a multidisciplinary team approach – we should all be passing on the same information.
- **Reiterate - talk to people living with HIV about the necessity of staying undetectable for U=U to be applicable.**
- Adherence management – discuss the importance of taking ART as prescribed to maintain health and prevent onward transmission.
- **Encourage engagement – ensure people living with HIV attend appointments and are fully engaged with their care and have knowledge of their viral load status. This means also discussing issues that may effect their adherence and engagement. What else is going on for them, life style, home, finances etc.**



THE NATIONAL HIV NURSES ASSOCIATION SUPPORTS

U = U

**THE RISK OF ACQUIRING HIV SEXUALLY FROM A PERSON WITH HIV WHO IS
UNDETECTABLE ON ANTIRETROVIRAL TREATMENT IS ZERO #UEQUALSU**

PREVENTIONACCESS.ORG

U=U*

UNDETECTABLE viral load means HIV IS UNTRANSMITTABLE

www.i-Base.info/u-equals-u



Michelle Ross,
clinIQ



Angelina Namiba,
Salamander Trust

“A person with sustained undetectable levels of HIV in their blood cannot transmit HIV to their sexual partners.”



Simon Collins,
HIV i-Base



Chloe Orkin,
Chair, British HIV Association
(BHIVA)



In PARTNER 1, couples (both straight and gay) had sex more than 58,000 times without condoms.

In PARTNER 2, in gay men only, couples had sex more than 77,000 times without condoms.

There were no linked transmissions in either study.

Other research includes the Rakai, HPTN 052 and Opposites Attract studies. It also includes the Swiss Statement that first published information about the risk being zero in 2008.

These and other studies are discussed in this online article.

The evidence for U=U: why negligible risk is zero risk

www.i-base.info/htb/32308

Does U=U apply to breastfeeding?

Although an undetectable viral load on ART also reduces the risk from breastfeeding, it doesn't reduce this risk to zero.

The have been cases where babies have become HIV positive from breastfeeding, even when the mother had an undetectable viral load.

Further information

The international U=U campaign raises awareness about this benefit from ART.

Currently, more than 720 organisations have joined from over 90 countries.

www.preventionaccess.org

i-Base has more information on U=U, including a longer version of this factsheet.

www.i-base.info/u-equals-u

U=U

Undetectable = Untransmittable

Did you know that having an undetectable viral load on HIV treatment (ART) stops HIV transmission?

ART is not only good for your health – it also protects your partners.

U=U means that you don't need to use condoms if you were only using them to stop HIV transmission.

Leading UK doctors and researchers strongly support the U=U statement.

“There should be no doubt that a person with sustained, undetectable levels of HIV in their blood cannot transmit HIV to their sexual partners.”



Professor Chloe Orkin,
Chair, British HIV Association

UK guidelines state that HIV doctors should talk to all their patients about how ART stops transmission.

What is U=U?

U=U stands for:

Undetectable = Untransmittable

It means that someone with an undetectable HIV viral load on ART cannot transmit HIV, even without using condoms or PREP.

What does U=U involve?

This protection from ART depends on:

- Taking ART every day.
- Having undetectable viral load for at least 6 months.
- Continuing to take your meds every day.

How can U=U not be a risk?

The quick answer is when HIV viral load is undetectable there is too little virus in sexual fluids for an infection to occur.

Produced by i-Base for Kobler@CWH (September 2018)

September 2018

www.i-Base.info

Any risk of sexual HIV transmission relates to viral load and being undetectable reduces this risk to zero.

Does U=U work with all HIV drugs?

Yes. Any ART that makes viral load undetectable will mean U=U.

Does U=U work for everyone?

Yes. The PARTNER studies included both gay and straight couples.

In straight couples sometimes the man was positive and in others the woman was positive.

Does U=U work for all types of sex?

Yes. The PARTNER studies collected information about the numbers of times people had oral, vaginal or anal sex. It also asked whether the negative partner was active or passive and whether there was ejaculation.

PARTNER reported zero transmissions for all situations.

Does this mean I can stop using condoms?

Whether you use condoms is a personal choice. Or a mutual choice with your partner.

Condoms reduce the risk of many STIs and they can effectively prevent pregnancy.

But if HIV is the only concern, then in the context of U=U, there is no reason to continue to use condoms.

My partner still wants to use condoms?

Whether your partner is HIV positive or HIV negative, they have to decide what is right for them.

They might want to use condoms for other reasons, or they might still worry about HIV transmission.

Sometimes it takes time for someone to accept new evidence, especially after using condoms for many years.

Do STIs affect U=U?

Based on the PARTNER study, U=U still works if, without realising it, one or both partners has an STI.

www.i-Base.info

It is important to monitor and treat STIs, but they are unlikely to affect U=U.

Is U=U widely accepted?

Yes, most leading HIV scientists and doctors now agree with the U=U statement.

This includes the British HIV Association (BHIVA), the International AIDS Society (IAS) and the US Center for Disease Control (CDC).

Will my doctor know about U=U?

Hopefully, yes. U=U has been headline news for at least two years. U=U is included in the BHIVA Standards for HIV Care (2018).

If your doctor doesn't tell you about U=U, then ask them.

How long does viral load need to be undetectable?

Guidelines recommend having an undetectable viral load for six months before relying on 100% protection from U=U.

This cautious approach is why guidelines refer to being on stable ART.

What if I miss my meds?

Missing your meds once will not change U=U.

But missing meds for 2-3 days might be enough with some combinations for viral load to become detectable.

Good adherence is essential for U=U.

What about viral load blips?

Sometimes viral load can “blip” just above 50 copies/mL.

Blips that stayed below 200 copies/mL did not affect the PARTNER results.

How do we know U=U is so effective?

Researchers have known for over 20 years that ART reduces all transmission risks. But it is only recently that the zero risk has been proved.

The PARTNER studies included couples where one partner was HIV positive on ART and the other HIV negative.

The positive partner needed to have an undetectable viral load and the couple needed to be having sex without condoms.

* Undetectable = Untransmittable

Poster produced by HIV i-Base for Kobler@CWH (July 2018)


September 2018

September 2018

Organisations endorsing BHIVA's position on U=U to date include:



I'll leave the final word to Laura Waters...



Resources

- <https://www.preventionaccess.org> – equal access to the HIV prevention revolution based on #ScienceNotStigma. All people living with HIV have a right to accurate and meaningful information about their social, sexual and reproductive health.
- <http://positiveseries.org> – offers downloadable and customizable social marketing campaigns to educate about U=U and encourage engagement in care.
- <http://i-base.info/u-equals-u/> - clear concise information.

References

- Cohen MS et al for the HPTN 052 Study Team. Prevention of HIV-1 infection with early antiretroviral therapy. Supplementary information. NEJM 2011; 365:493-505. <http://www.nejm.org/doi/full/10.1056/NEJMoa1105243>
- Cohen MS et al. Final results of the HPTN 052 randomized controlled trial: antiretroviral therapy prevents HIV transmission. IAS 2015, 19 – 22 July 2015, Vancouver. MOAC0101LB. <http://dx.doi.org/10.7448/IAS.18.5.20482>
- Fox J et al. Quantifying sexual exposure to HIV within an HIV-serodiscordant relationship: development of an algorithm. AIDS 2011, 25:1065–1082. DOI:10.1097/QAD.0b013e328344fe4a. Free online access. http://journals.lww.com/aidsonline/Abstract/2011/05150/Quantifying_sexual_exposure_to_HIV_within_an.7.aspx
- Grulich A et al. HIV treatment prevents HIV transmission in male serodiscordant couples in Australia, Thailand and Brazil. IAS 2017, Paris. Oral abstract TUAC0506LB. <http://programme.ias2017.org/Abstract/Abstract/5469>
- Rodger A et al. HIV transmission risk through condomless sex if HIV+ partner on suppressive ART: PARTNER Study. 21st CROI, 3-6 March 2014, Boston. Oral late breaker abstract 153LB. <http://www.croiwebcasts.org/console/player/22072> (webcast)
- Rodger AJ et al for the PARTNER study group. Sexual activity without condoms and risk of HIV transmission in serodifferent couples when the HIV-positive partner is using suppressive antiretroviral therapy. JAMA, 2016;316(2):1-11. DOI: 10.1001/jama.2016.5148. (12 July 2016). Full free access. <http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2016.5148>
- Vernazza P et al. HIV-positive individuals not suffering from any other STD and adhering to an effective antiretroviral treatment do not transmit HIV sexually. (Les personnes séropositives ne souffrant d'aucune autre MST et suivant un traitement antirétroviral efficace ne transmettent pas le VIH par voie sexuelle). Bulletin des médecins suisses 89 (5), 30 January 2008. Included with English translation. http://i-base.info/qa/wp-content/uploads/2008/02/Swiss-Commission-statement_May-2008_translation-EN.pdf (PDF)